

LEVEL

MD A 0 75577

ON THE CANDIDATE PROBLEM WITH A RANDOM NUMBER OF CANDIDATES

by

C. DERMAN, G. J. LIEBERMAN S. ROSS

TECHNICAL REPORT NO. 192

AUGUST 1979

(4) TR-192

SUPPORTED BY THE ARMY AND NAVY
UNDER CONTRACT NOO014-75-C-0561 (NR-047-200)
WITH THE OFFICE OF NAVAL RESEARCH

Gerald J. Lieberman, Project Director

Reproduction in Whole or in Part is Permitted for any Purpose of the United States Government Approved for public release; distribution unlimited

Accession For

FTIS GRAAI

IDC TAB

Unammounced

Justification

By


Distribution/

Availability Codes

Availand/or

Special

DEPARTMENT OF OPERATIONS RESEARCH AND DEPARTMENT OF STATISTICS STANFORD UNIVERSITY STANFORD, CALIFORNIA



402 766

ON THE CANDIDATE PROBLEM WITH A RANDOM NUMBER OF CANDIDATES 1

by

C. Derman, Columbia University G.J. Lieberman, Stanford University S.M. Ross, University of California, Berkeley

1. Introduction

In the problem under consideration a decision maker has a total of M candidates to interview sequentially. The decision maker must either accept or reject the candidate being interviewed after he has been ranked with respect to his predecessors. Once rejected a candidate cannot be reconsidered; once a candidate is accepted no further interviews are carried out. The objective is to select the candidate in such a way as to maximize the probability of choosing the best of all M candidates (assuming every ordering of interviews is equally likely).

In the classical version, M = m is fixed and the optimal selection policy is to interview r - 1 candidates without choosing any; then select the first leading candidate interviewed thereafter. Here

$$r = r^*(m) = min \left\{ r \ge 1 \middle| \sum_{k=r}^{m-1} 1/k \le 1 \right\}.$$

This research was supported in part by the office of Naval Research under Contracts N00014-75-C-0620 with Columbia University, N00014-75-C-0561 with Stanford University, and N00014-77-C-0299 with the University of California, by the Air Force Office of Scientific Research (AFSC), USAF, under Grant AFOSR-77-3213 with the University of California, and by the National Science Foundation under Grant MCS-7825-146 with Columbia University.

(See Chow, et.al. [1] p. 51 or Derman [3] p. 118.)

Rasmussen [5] and Rasmussen and Robbins [6] generalize the problem to the case where M is random with known probability function $P(M = m) = p_m$, $m = 1, ..., N, N < \infty$. The approach taken in [6] rests on the presumption (asserted in [5]) that the optimal solution policy possesses the above simple form and that one need only find the value of r that maximizes

$$\varphi(r) = \begin{cases} \sum_{m=1}^{N} p_m/m, & \text{if } r = 1 \\ N & \sum_{m=1}^{m-1} 1/k, & \text{if } r \ge 2 \end{cases},$$

the a priori probability of selecting the best candidate under the rule which lets the first r-1 go by and then selects the first leader that occurs thereafter.

The assertion of [5], however, is not true for every probability function $\{p_m\}$. In Section 2 a counter-example shows this. In Section 3 a sufficient condition is given for the optimal policy to have the simple form. The device for generating the condition is the "one-step look-ahead" criterion given by Derman and Sacks [4] (also by Chow and Robbins [2]). It turns out that this criterion applies if and only if $\varphi(r)$ is unimodal. The condition given here for the criterion to apply improves on the one given in [6] for $\varphi(r)$ to be unimodal.

2. Counter-example

If $\{p_m\}$ were a two-point distribution with large probability mass at M=2 (say) and small, but, positive, probability mass at M=N(N | large), a simple selection policy would have r=1 or 2. Intuitively, however, it would seem that a better policy would be to select the second candidate if he was better than the first and if not to let a number of candidates go by before deciding to choose the next leading candidate. Formally, suppose

$$p_2 = 1 - \epsilon$$
, $p_N = \epsilon$.

Then

$$\varphi(1) = \frac{1-\epsilon}{2} + \epsilon/N$$

$$\varphi(2) = \frac{1-\epsilon}{2} + \frac{\epsilon}{N} \sum_{k=1}^{N-1} 1/k$$

$$\varphi(\mathbf{r}) < \epsilon, 3 \le \mathbf{r} \le \mathbf{N}$$
.

So for N > 2 and E small

$$\varphi(2) = \max_{\mathbf{r}} \varphi(\mathbf{r})$$

However, consider the policy that stops at r=2, if possible, or, if M=N, at the first leading candidate after i-1 candidates have been interviewed, where i shall be appropriately determined. Let φ denote the probability of selecting the best candidate under this policy. Then,

$$\varphi = P\{M = 2\} \quad P\{\text{Selecting Best Candidate} | M = 2\}$$

$$+ P\{M = N\} \quad P\{\text{Selecting Best Candidate} | M = N\}$$

$$= \frac{1-\epsilon}{2} + \epsilon \left\{ 1/2 \cdot 2/N + 1/2 \cdot \frac{N-2}{N} \left(\frac{i-3}{N-2} \sum_{k=i-3}^{N-3} 1/k \right) \right\}$$

$$= \frac{1-\epsilon}{2} + \epsilon/N \left\{ 1 + \frac{i-3}{2} \sum_{k=i-3}^{N-3} 1/k \right\} .$$

Thus.

$$\varphi > \varphi(2)$$

if

$$1 + \frac{i-3}{2} \sum_{k=i-3}^{N-3} 1/k \ge \sum_{k=1}^{N-1} 1/k$$
.

If i is any integer greater than 5 and N is sufficiently large, the latter inequality will hold.

3. One-step Look Ahead Policies

Consistent with our use of the expression in the foregoing, we say a leading candidate is being interviewed if he ranks above all previously interviewed candidates. A <u>one-step look ahead policy</u> is one that accepts a leading candidate if and only if the probability that he is the best exceeds the probability that one more leader will be interviewed and that he will be the best. Applying a theorem by Derman and Sacks [4] (also Chow and Robbins [2]), the one-step look ahead criterion yields an optimal selection policy if there is 30 - 1 such that one-step look ahead accepts (rejects) a leader among the first j interviewed if $j > r - 1(j \le r - 1)$.

Let

$$p(m|i) = \frac{p_m}{\sum_{k \ge i} p_k}, m \ge i,$$

denote the conditional probability that M = m given $M \ge i$. Let \mathscr{E}_j denote the event that the j^{th} candidate interviewed is a leader, \mathscr{E}_i denote the event that the j^{th} candidate is the best. Then

$$p\{\mathscr{F}_{j} | M \ge j, \mathscr{E}_{j}\} = j \sum_{m=j}^{N} \frac{p(m|j)}{m} \equiv R(j)$$
 (say).

Let \mathcal{G}_j denote the event that the jth candidate observed is a leader, one more leader will be observed and that he will be the best. Then

$$P\{\mathscr{S}_{j} | M \geq j, \mathscr{E}_{j}\} = \sum_{m=j+1}^{N} p(m|j) \sum_{k=j+1}^{m} \frac{j}{k(k-1)} \frac{k}{m}$$

$$= j \sum_{m=j+1}^{N} \frac{p(m|j)}{m} \sum_{k=j}^{m-1} 1/k$$

$$\equiv S(j) \qquad (say) \qquad .$$

Thus, the one-step look policy is optimal if for some r - 1

(1)
$$R(j) - S(j) < 0, j \le r - 1$$

 $\ge 0, j \ge r$.

Now, interpreting $\sum_{k=j}^{j-1} 1/k = 0,$

$$R(j) - S(j) = j \left\{ \sum_{m=j}^{N} \frac{p(m|j)}{m} - \sum_{m=j+1}^{N} \frac{p(m|j)}{m} \sum_{k=j}^{m-1} 1/k \right\}$$

$$= \frac{1}{N} \left\{ \sum_{m=j}^{N} \frac{p_m}{m} \left(1 - \sum_{k=j}^{m-1} 1/k \right) \right\}$$

$$\begin{array}{ccc}
 & & & \downarrow \\
 & & & & \downarrow \\
 & & \downarrow \\$$

where $\psi(j)$ is the expression within the last brackets. The sign of R(j) - S(j) is the same as that of $\psi(j)$. Thus, from (1) follows

Proposition 1: A one-step look ahead policy is optimal if for some r - 1

(2)
$$\psi(j) \leq 0, \ j = 1, \dots, \ r-1$$
 $\psi(j) > 0, \ j = r, \dots, \ N$.

The hypothesis of proposition 1 is equivalent to $\varphi(\mathbf{r})$ being unimodal since, as given in [6],

$$\varphi(j) - \varphi(j+1) - \psi(j), j - 1, ..., N - 1$$
.

Now let

$$H_j = P_j - \sum_{m=j+1}^{N} P_m/m, j = 1, ..., N-1$$
.

In terms of H, we state

Condition (*): $H_{i} \geq 0$ implies $H_{i} \geq 0$ for $i \geq j$.

We now have

Theorem 1: If condition (*) holds, then there is an r such that

(2) holds (i.e., a one-stop look ahead policy is optimal).

Proof: In Rasmussen and Robbins [6] it is given that

$$\psi(j) - \psi(j+1) = H_j/j$$
, $j = 1, ..., N-1$.

Let i be any integer such that

 $\psi(i) \geq 0$.

If $H_i < 0$, then

$$\psi(i+1) \geq \psi(i)$$

> 0 .

If $H_i \ge 0$, then by condition (*)

$$H_j \ge 0$$
, j = i + 1, ..., N - 1,

and hence, $\psi(j)$ is non-increasing in j for all j, $i \leq j \leq N$. However,

 $\psi(N) \ge 0;$

therefore, $\psi(j) \ge 0$ for all j, $i \le j \le N$. Thus, in both cases $\psi(i+1) \ge 0$, proving the theorem.

Remarks: Condition (*) is weaker than the condition $p_1 \leq p_2 \leq \cdots \leq p_N$ given in [6]. It is also satisfied by the geometric distribution over 1, ..., N, a distribution for which $\varphi(r)$ is unimodal (as stated in [6]). It is not necessary to restrict N to being finite; under condition (*) Theorem 1 will hold with N = ∞ since, in this case, $\lim_{j \to \infty} \psi(j) = 0$.

REFERENCES

- [1] Chow, Y.S., H. Robbins, and D. Siegmund (1971), Great Expectations: The Theory of Optimal Stopping, Houghton Mifflin Company, Boston.
- [2] Chow, Y.S., and H. Robbins (1961), A Martingale System Theorem Applications, Proc. Berkeley Symposium, Math Stat. Prob., 4th., 93-104.
- [3] Derman, C. (1970), Finite State Markovian Decision Processes, Academic Press, New York.
- [4] Derman, C. and J. Sacks (1960), Replacement of Periodically Inspected Equipment (An Optimal Stopping Rule), Naval Research Logistics Quarterly, 7, 597-607.
- [5] Rasmussen, W.T. (1975), A Generalized Choice Problem, J. Optimization Theory Appl., 15, 311-325.
- [6] Rasmussen, W.T., and H. Robbins (1975), The Candidate Problem with Unknown Population Size, J. Appl. Prob., 12, 692-701.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE /When Dete !	Entered)	
REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	1. RECIPIENT'S CATALOG NUMBER
ON THE CANDIDATE PROBLEM WITH A RANDOM NUMBER OF CANDIDATES		S. TYPE OF REPORT & PERIOD COVERED Technical Report 6. PERFORMING ORG. REPORT NUMBER
7. Author(*) C. Derman, G.J. Lieberman and S. Ross		N00014-75-C-0561
 PERFORMING ORGANIZATION NAME AND ADDRESS Department of Operations Research and Department of Statistics Stanford University, Stanford, California 94305 		16. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT HUMBERS (NR-047-200)
Operations Research, Code 434 Office of Naval Research		August 10, 1979
Arlington, Virginia 22217 14. MONITORING AGENCY NAME & ADDRESS/II different from Controlling Office)		10 18. SECURITY CLASS. (et this report) Unclassified 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
APPROVED FOR PUBLIC RELEASE: DISTR	RIBUTION IS UNLIM	ITED

17. DISTRIBUTION STATEMENT (of the obstract entered in Block 30, if different from Report)

18. SUPPLEMENTARY NOTES

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

Candidate problem One-step look-ahead policy Choice problem Random number of candidates

10. ABSTRACT (Continue on reverse elde il necessory and identity by bleck number)

A decision maker has a total of M candidates to interview sequentially. The decision maker must either accept or reject the candidate being interviewed after he has been ranked with respect to his predecessors. Once rejected a candidate cannot be reconsidered; once a candidate is accepted no further interviews are carried out. The objective is to select the candidate in such a way as to maximize the probability of choosing the best of all M candidates (assuming every ordering of interviews is equally likely). ..

(Continued)

In the classical version, M = m is fixed and the optimal selection policy is to interview r-1 candidates without choosing any; then select the first leading candidate interviewed thereafter. Here

$$r = r^{*}(m) = \min \left\{ r \geq 1 \middle| \sum_{k=r}^{m-1} 1/k \leq 1 \right\}.$$

Rasmussen (1975) and Rasmussen and Robbins (1975) generalize the problem to the case where M is random with known probability function $P(M=m)=p_m$, $m=1,\ldots,N,N<\infty$. The approach taken by Rasmussen and Robbins rests on the presumption (asserted by Rasmussen) that the optimal solution policy possesses the above simple form and that one need only find the value of r that maximizes

$$\varphi(r) = \begin{cases} \sum_{m=1}^{N} p_m/m, & \text{if } r = 1 \\ N & m-1 \\ (r-1) \sum_{m=r}^{N} p_m/m & \sum_{k=r-1}^{m-1} 1/k, & \text{if } r \ge 2 \end{cases},$$

the a priori probability of selecting the best candidate under the rule which lets the first r-1 go by and then selects the first leader that occurs thereafter.

Rasmussen's result, however, is not true for every probability function $\{p_m\}$. In Section 2 a counter-example shows this. In Section 3 a sufficient condition is given for the optimal policy to have the simple form. The device for generating the condition is the "one-step look-ahead" criterion given by Derman and Sacks (1960) (also by Chow and Robbins (1961)). It turns out that this criterion applies if and only if $\varphi(r)$ is unimodal. The condition given here for the criterion to apply improves on the one given by Rasmussen and Robbins (1975) for $\varphi(r)$ to be unimodal.

